

## Focus on topological quantum computation

**Jiannis K. Pachos**

School of Physics and Astronomy, University of Leeds, Leeds, LS2 9JT,  
United Kingdom

E-mail: [j.k.pachos@leeds.ac.uk](mailto:j.k.pachos@leeds.ac.uk)

**Steven H. Simon**

Rudolf Peierls Centre for Theoretical Physics, University of Oxford, OX1 3NP,  
United Kingdom

E-mail: [s.simon1@physics.ox.ac.uk](mailto:s.simon1@physics.ox.ac.uk)

**Abstract.** Topological quantum computation started as a niche area of research aimed at employing particles with exotic statistics, called anyons, for performing quantum computation. Soon it evolved to include a wide variety of disciplines. Advances in the understanding of anyon properties inspired new quantum algorithms and helped in the characterisation of topological phases of matter and their experimental realisation. The conceptual appeal of topological systems as well as their promise for building fault-tolerant quantum technologies fuelled the fascination in this field. This ‘focus on’ brings together several of the latest developments in the field and facilitates the synergy between different approaches.

The great promise of quantum computers has to be balanced against the great difficulty of actually building them. Foremost among the difficulties is the fundamental challenge of defeating decoherence and errors. Small improvements to current strategies may not be sufficient to overcome this problem; radically new ideas may be required. Topological quantum computation is precisely such a new and different approach [1]. It employs many-body physical systems with the unique property of encoding and processing quantum information in a naturally fault-tolerant way.

Research on topological quantum computation has become a highly interdisciplinary field, with frontiers in physics, mathematics, and computer science. Moreover, advances in the theoretical understanding of abstract topology [2], in physical realisations of topological matter [3], and in computational paradigms [4] have been closely interrelated, with developments in one area strongly influencing the others. Recent years have witnessed significant theoretical and experimental developments. These include major experimental and theoretical advances in fractional quantum Hall systems that support the existence of non-Abelian anyons [5] – the building block of topological quantum computation – as well as the predication and experimental discovery of novel spin-orbit systems such as topological insulators [6].

The aim of this focus issue is to bring together the latest developments in a variety of areas with the goal of promoting new results to a wider community of scientists and advancing the synergy between different approaches. To a large extent we believe this has been a success.

One of the dominant directions of the field has been towards better understanding topological matter by investigating phase transitions between topological phases — how one phase condenses out of another, often in some sort of confinement transition. Loop models, as the simplest of all topological phases, were studied by Monte-Carlo methods [7]. Semi-analytic high order perturbation theory methods were applied to study transitions in  $\mathbb{Z}_N$  topological phases [8]. In some cases powerful purely analytic approaches could be applied to study phase transitions in detail [9, 10, 11]. There was substantial focus on identifying order parameters for such topological phase transitions to try to draw analogies with the conventional Landau theory of phase transitions [12, 13]. Despite this progress, and although some general principles are now established [14], a unified theory of the details of phase transitions between topological phases is still absent and remains an active topic of research.

Another emerging direction in the field has been the study of Majorana fermions and the physical systems that may harbour them. In the past few years, several proposals were made for creating analogs of chiral p-wave superconductors using spin-orbit coupled superconductors, which would host Majorana zero modes at defects such as vortices. Braiding these defects would result in non-Abelian operations on a degenerate ground state [15] (see [16] for a review of these ideas). Indeed, this development became such an active field that it called for a separate focus issue [17]. However, before that issue was created many groundbreaking works on the topic ended up in the current focus issue. One direction is in determining the parameters necessary for obtaining topological superconductivity in spin orbit coupled systems [18]. Another direction is the proposal of experiments (in this case Ettingshausen effect experiments [19] or quantum point contact tunnelling experiments [20]) that might establish the existence of Majorana modes in these systems. Yet another direction is in the development of device designs to read out [21, 22] or manipulate [23] such Majoranas qubits. Majoranas, as perhaps the simplest non-Abelian object, were also studied in several other contexts including phase transitions [9], fractional quantum Hall effect [24], the Kitaev Honeycomb model [25, 26, 27, 28], Hanbury-Brown-Twiss experiments, [29] and quench dynamics of spin chains [30]. It is certain that the study of Majorana physics will be a main direction of the field in the future.

A perennially dominant direction since the inception of the field has been the study of fractional quantum Hall effects [31, 3]. This is perhaps not surprising being that quantum Hall effect in semiconductors remains the only physical system convincingly established as nontrivial topological matter. In this issue quantum Hall physics was also discussed in other physical realisations ranging from mono-layer and bilayer graphene [32] to cold atoms in non-Abelian gauge fields [33, 34]. The presence of extra degrees of freedom, such as spin and valley, over the conventionally studied spinless electrons, gives the freedom to have richer physics, including charged spin textures [35]. In another direction, a nice theoretical advance (the extension of a many year project by some of the same authors and others) was the development of a conformal field theory approach to the quantum Hall hierarchies [36]. As in the broader field of condensed matter, understanding the patterns of entanglement of fractional Hall states has becoming an increasingly important and interesting topic [37, 38].

While several experimental works have recently been performed attempting to demonstrate non-Abelian quasiparticles in the fractional quantum Hall effect [39, 40], the interpretation of these experiments is very controversial, and this remains a forefront of research. Several works in the current issue were aimed at either providing

other methods to make this demonstration, for example by measuring tunnelling spectra [41], or were aimed to examine additional physics of these experiments (in this case the Zeno effect [24], or disorder in quasiparticle lattice [42]) not previously considered in predictions. In our entire focus issue, disappointingly, only a single submitted manuscript was actually a real experiment [43], and this, although a nice experiment demonstrating Aharonov-Bohm oscillations in quantum Hall samples, remains quite a distance from the desired result that would convincingly show non-Abelian physics.

Back to the theoretical front we note that perhaps one of the most studied models has been the Kitaev honeycomb model [25]. Unsurprisingly this continued to provide fertile ground for further thought. Within this framework the interactions between non-Abelian vortices and the nucleation of new topological phases was studied [26] and the braiding statistics were evaluated explicitly to very high accuracy [27]. A square-octagon variant of the Kitaev honeycomb model was shown to exhibit a rather remarkably rich phase diagram [28].

Another of the most celebrated models of topological matter is the toric code model [1]. In one work, the thermal stability of this model was examined [44]. Another work considers how this model might actually be simulated using trapped ions [45]. Since its inception, the toric code model has since been generalised to  $\mathbb{Z}_N$  models [8], non-Abelian discrete gauge theories [10], and finally to their most general form, the Levin-Wen models [46, 9, 47, 11, 48]. One work in this issue geometrically reinterpreted the Levin-Wen partition functions as knot invariants [47]. Another work explores (at least in one dimension) generalising to theories based on non-unitary conformal field theories [48].

On the more computational side of research, one of the key ideas that launched the field was the realisation that evaluation of the Jones polynomial, a “hard” problem for a classical computer, might not be “hard” for an anyon computer [49]. Indeed, evaluation of Jones polynomials [4, 50] is perhaps one of the most natural algorithms for a topological quantum computer. In this issue, extensions of this line of thought have resulted in advances in determining the precise complexity of this quantum algorithm [51] and in its relationship to Khovanov homology — a natural “categorification” of the Jones polynomial [52].

The quantum gates produced by anyonic braiding are very specific, tightly connected to the statistical properties of the anyons. In this focus issue, it was shown that no computationally universal anyon system can be built where two qubit gates do not have finite leakage errors [53]. This further draws attention towards the necessity of approximating quantum gates to high accuracy (even though they must remain imperfect). One approach pursued towards designing such high accuracy gates is to use hashing techniques [54]. Finally, ideas from cluster state quantum computation have been applied to the realm of topological models [55, 56].

As witnessed from these contributions it is clear that the field of topological quantum computing, and topological matter in general, has continued to progress at a remarkable rate. Despite these advances, several fundamental questions still remain open. Arguably, two main directions can be identified: building systems where non-Abelian statistics can be conclusively measured and finding a topologically inspired quantum memory that can successfully combat physically induced errors. It is the hope and vision of the editors that bringing together methods and techniques from wide range of fields can eventually enable the construction of topological quantum computation.

## References

- [1] Kitaev A Yu 2003 Fault-tolerant quantum computation by anyons, *Annals Phys.* **303** 2-30
- [2] Jones V F R 1985 A polynomial invariant for knots via von Neumann algebras *Bull. Amer. Math. Soc.* **12** 103
- [3] Nayak C, Simon S H, Stern A, Freedman M, and Das Sarma S 2008 Non-Abelian Anyons and Topological Quantum Computation *Rev. Mod. Phys.* **80** 1083
- [4] Freedman M, Kitaev A, Larsen M J and Wang Z 2003 Topological quantum computation *Bull. Amer. Math. Soc.* **40** 31
- [5] Moore G and Read N 1991 Nonabelions in the fractional quantum Hall effect, *Nucl. Phys. B* **360**, 362
- [6] Hasan M Z and Kane C L 2010 Topological Insulators *Rev. Mod. Phys.* **82** 3045
- [7] Herdman C M and Whaley K B 2011 Loop condensation in the triangular lattice quantum dimer model *New J. Phys.* **13** 085001
- [8] Schulz M D, Dusuel S, Orus R, Vidal J and Schmidt K P 2012 Breakdown of a perturbed  $\mathbb{Z}_N$  topological phases *New J. Phys.* **14** 025005
- [9] Burnell F J, Simon S H and Slingerland J K 2012 Phase transitions in topological lattice models via topological symmetry breaking *New J. Phys.* **14** 015004
- [10] Bombin H and Martin-Delgado M A 2011 Nested topological order *New J. Phys.* **13** 125001
- [11] Ludwig A W W, Poilblanc D, Trebst S and Troyer M 2011 Two-dimensional quantum liquids from interacting non-Abelian anyons *New J. Phys.* **13** 045014
- [12] Bais F A and Romers J C 2012 The modular S-matrix as order parameter for topological phase transitions *New J. Phys.* **14** 035024
- [13] Gregor K, Huse D A, Moessner R and Sondhi S L 2011 Diagnosing deconfinement and topological order *New J. Phys.* **13** 025009
- [14] Bais F A and Slingerland J K 2009 Condensate induced transitions between topologically ordered phases *Phys. Rev. B* **79** 045316
- [15] Read N 2012 Topological phases and quasiparticle braiding, *Physics Today* **65**, 38
- [16] Alicea J 2012 New directions in the pursuit of Majorana fermions in solid state systems *Reports on Progress in Physics* **75** 076501
- [17] Fu L and Marcus C M Focus Issue on Majorana Fermions in Condensed Matter *New Journal of Physics*
- [18] Tewari S, Stanescu T D, Sau J D and Das Sarma S 2011 Topologically non-trivial superconductivity in spin-orbit-coupled systems: bulk phases and quantum phase transitions *New J. Phys.* **13** 065004
- [19] Hou C-Y, Shtengel K, Refael G and Goldbart P M 2012 Ettingshausen effect due to Majorana modes *New J. Phys.* **14** 105005
- [20] Wimmer M, Akhmerov A R, Dahlhaus J P and Beenakker C W J 2011 Quantum point contact as a probe of a topological superconductor *New J. Phys.* **13** 053016
- [21] Hassler F, Akhmerov A R and Beenakker C W J 2011 The top-transmon: a hybrid superconducting qubit for parity-protected quantum computation *New J. Phys.* **13** 095004
- [22] Hassler F, Akhmerov A R, Hou C-Y and Beenakker C W J 2010 Anyonic interferometry without anyons: how a flux qubit can read out a topological qubit *New J. Phys.* **12** 125002
- [23] van Heck B, Akhmerov A R, Hassler F, Burrello M and Beenakker C W J 2012 Coulomb-assisted braiding of Majorana fermions in a Josephson junction array *New J. Phys.* **14** 035019
- [24] Clarke D J and Shtengel K 2011 Edge-induced qubit polarization in systems with Ising anyons *New J. Phys.* **13** 055005
- [25] Kitaev A 2006 Anyons in an exactly solved model and beyond *Annals of Physics* **321** 2–111
- [26] Lahtinen V 2011 Interacting non-Abelian anyons as Majorana fermions in the honeycomb lattice model *New J. Phys.* **13** 075009
- [27] Bolukbasi A T and Vala J 2012 Rigorous calculations of non-Abelian statistics in the Kitaev honeycomb model *New J. Phys.* **14** 045007
- [28] Kells G, Kailasvuori J, Slingerland J K and Vala J 2011 Kaleidoscope of topological phases with multiple Majorana species *New J. Phys.* **13** 095014
- [29] Bose S and Sodano P 2011 Nonlocal Hanbury-Brown-Twiss interferometry and entanglement generation from Majorana bound states *New J. Phys.* **13** 085002
- [30] DeGottardi W, Sen D and Vishveshwara S 2011 Topological phases, Majorana modes and quench dynamics in a spin ladder system *New J. Phys.* **13** 065028
- [31] Stormer H L 1999 Nobel Lecture: The fractional quantum Hall effect *Reviews of Modern Physics* **71** 875
- [32] Abanin D A, Papić Z, Barlas Y and Bhatt R N 2012 Stability of the  $k=3$  Read-Rezayi state in

- chiral two-dimensional systems with tunable interactions *New J. Phys.* **14** 025009
- [33] Palmer R N and Pachos J K 2011 Fractional quantum Hall effect in a  $U(1) \times SU(2)$  gauge field *New J. Phys.* **13** 065002
- [34] Estienne B, Haaker S M and Schoutens K 2011 Particles in non-Abelian gauge potentials: Landau problem and insertion of non-Abelian flux *New J. Phys.* **13** 045012
- [35] Romers J, Huijse L and Schoutens K 2011 Charged spin textures over the Moore-Read quantum Hall state *New J. Phys.* **13** 045013
- [36] A general approach to quantum Hall hierarchies J Suorsa, S Viefers and T H Hansson 2011 *New J. Phys.* **13** 075006
- [37] Sterdyniak A, Bernevig B A, Regnault N and Haldane F D M 2011 The hierarchical structure in the orbital entanglement spectrum of fractional quantum Hall systems *New J. Phys.* **13** 105001
- [38] Friedman B A, Levine G C and Luna D 2011 Entanglement entropy of random fractional quantum Hall systems *New J. Phys.* **13** 055006
- [39] Willett R L, Nayak C, Shtengel K, Pfeiffer L N and West K W 2013 Magnetic-Field-Tuned Aharonov-Bohm Oscillations and Evidence for Non-Abelian Anyons at  $\nu = 5/2$  *Phys. Rev. Lett.* **111** 186401
- [40] arXiv:1112.3400 Braiding of Abelian and Non-Abelian Anyons in the Fractional Quantum Hall Effect, Sanghun An, P. Jiang, H. Choi, W. Kang, S. H. Simon, L. N. Pfeiffer, K. W. West, K. W. Baldwin
- [41] Hu Z-X, Lee K H, Rezayi E H, Wan X and Yang K 2011 Scaling and non-Abelian signature in fractional quantum Hall quasiparticle tunneling amplitude *New J. Phys.* **13** 035020
- [42] Kraus Y E and Stern A 2011 Majorana fermions on a disordered triangular lattice *New J. Phys.* **13** 105006
- [43] Choi H, Jiang P, Godfrey M D, Kang W, Simon S H, Pfeiffer L N, West K W and Baldwin K W 2011 Aharonov-Bohm-like oscillations in Fabry-Perot interferometers *New J. Phys.* **13** 055007
- [44] Al-Shimary A, Wootton J R and Pachos J K 2013 Lifetime of topological quantum memories in thermal environment *New J. Phys.* **15** 025027
- [45] Müller M, Hammerer K, Zhou Y L, Roos C F and Zoller P 2011 Simulating open quantum systems: from many-body interactions to stabilizer pumping *New J. Phys.* **13** 085007
- [46] Levin M A and Wen X-G 2005 String-net condensation: A physical mechanism for topological phases *Phys. Rev. B* **71** 045110
- [47] Burnell F J and Simon S H 2011 A Wilson line picture of the Levin-Wen partition functions *New J. Phys.* **13** 065001
- [48] Ardonne E, Gukelberger J, Ludwig A W W, Trebst S and Troyer M 2011 Microscopic models of interacting Yang-Lee anyons *New J. Phys.* **13** 045006
- [49] Freedman M H 1998 P/NP and the quantum field computer *Proc. Natl. Acad. Sci.* **95** 98
- [50] Aharonov D, Jones V and Landau Z 2009 A Polynomial Quantum Algorithm for Approximating the Jones Polynomial *Algorithmica* **55** 395
- [51] Aharonov D and Arad I 2011 The BQP-hardness of approximating the Jones polynomial *New J. Phys.* **13** 035019
- [52] Kauffman L H 2011 Topological quantum information, virtual Jones polynomials and Khovanov homology *New J. Phys.* **13** 125007
- [53] Ainsworth R and Slingerland J K 2011 Topological qubit design and leakage *New J. Phys.* **13** 065030
- [54] Burrello M, Mussardo G and Wan X 2011 Topological quantum gate construction by iterative pseudogroup hashing *New J. Phys.* **13** 025023
- [55] Horsman C 2011 Quantum pictorialism for topological cluster-state computing *New J. Phys.* **13** 095011
- [56] Brown B J, Son W, Kraus C V, Fazio R and Vedral V 2011 Generating topological order from a two-dimensional cluster state using a duality mapping *New J. Phys.* **13** 065010